

The type locality of *humboldti* is considerably east of the known distribution of any species with similar genitalia. The other species are from California and almost without exception from the coastal region of southern California.

REFERENCES

- DAWSON, R. W. 1919. New species of *Serica* (Scarabaeidae).-I. J. New York Ent. Soc. 27:29-32.
 _____ 1919. New species of *Serica* (Scarabaeidae).-II. J. New York Ent. Soc. 27:223-224.
 _____ 1920. New species of *Serica* (Scarabaeidae). III. J. New York Ent. Soc. 28:208-211.
 _____ 1921. New species of *Serica* (Scarabaeidae). IV. J. New York Ent. Soc. 29:160-168.
 _____ 1922. New species of *Serica* (Scarabaeidae). V. J. New York Ent. Soc. 30:154-169.
 _____ 1932. New species of *Serica* (Scarabaeidae). VI. J. New York Ent. Soc. 40:529-548.
 _____ 1933. New species of *Serica* (Scarabaeidae). VII. J. New York Ent. Soc. 41:435-440.
 _____ 1947. New species of *Serica* (Scarabaeidae). VIII. J. New York Ent. Soc. 55:223-235.
 _____ 1952. New species of *Serica* (Scarabaeidae). IX. J. New York Ent. Soc. 60:65-77.
 _____ 1967. New and little known species of *Serica* (Coleoptera: Scarabaeidae). X. J. New York Ent. Soc. 75:161-178.



PSYDRUS PICEUS LeCONTE FROM ARIZONA
(COLEOPTERA: CARABIDAE)

To my knowledge the following are the first published records of *Psydrus piceus* Lec. from Arizona: Cochise County, Chiricahua Mountains, Rustler's Park, 7-VIII-48, G. E. Ball, 8500' (1); Apache County, 3 miles north of Alpine on U. S. Highway 666, 8-9-VIII-69, G. E. and K. E. Ball, 8100' (3); Navaho County, Pinetop, 3-VIII-73, Scott McCleve, 7000' (1). All 5 specimens were found under the bark of pine logs. I thank Dr. Ball for permission to include and publish his records with mine.—Scott McCleve, 2210 13th Street, Douglas, AZ 85607

